

Comparison of lignin peroxidase, horseradish peroxidase and laccase in the oxidation of methoxybenzenes

Philip J. KERSTEN,*|| B. KALYANARAMAN,† Kenneth E. HAMMEL,‡ Bengt REINHAMMAR§ and T. Kent KIRK*¶

*Forest Products Laboratory, USDA Forest Service, Madison, WI 53705 and Department of Bacteriology, University of Wisconsin, Madison, WI 53706, †National Biomedical ESR Center, Medical College of Wisconsin, Milwaukee, WI 53226, ‡Department of Chemistry, SUNY College of Environmental Sciences and Forestry, Syracuse, NY 13210, U.S.A., and §Department of Biochemistry and Biophysics, Chalmers University of Technology, Göteborg, Sweden

Lignin peroxidase oxidizes non-phenolic substrates by one electron to give aryl-cation-radical intermediates, which react further to give a variety of products. The present study investigated the possibility that other peroxidative and oxidative enzymes known to catalyse one-electron oxidations may also oxidize non-phenolics to cation-radical intermediates and that this ability is related to the redox potential of the substrate. Lignin peroxidase from the fungus *Phanerochaete chrysosporium*, horseradish peroxidase (HRP) and laccase from the fungus *Trametes versicolor* were chosen for investigation with methoxybenzenes as a homologous series of substrates. The twelve methoxybenzene congeners have known half-wave potentials that differ by as much as ~ 1 V. Lignin peroxidase oxidized the ten with the lowest half-wave potentials, whereas HRP oxidized the four lowest and laccase oxidized only 1,2,4,5-tetramethoxybenzene, the lowest. E.s.r. spectroscopy showed that this congener is oxidized to its cation radical by all three enzymes. Oxidation in each case gave the same products: 2,5-dimethoxy-*p*-benzoquinone and 4,5-dimethoxy-*o*-benzoquinone, in a 4:1 ratio, plus 2 mol of methanol for each 1 mol of substrate. Using HRP-catalysed oxidation, we showed that the quinone oxygen atoms are derived from water. We conclude that the three enzymes affect their substrates similarly, and that whether an aromatic compound is a substrate depends in large part on its redox potential. Furthermore, oxidized lignin peroxidase is clearly a stronger oxidant than oxidized HRP or laccase. Determination of the enzyme kinetic parameters for the methoxybenzene oxidations demonstrated further differences among the enzymes.

INTRODUCTION

Growing evidence indicates that the lignin peroxidase of the fungus *Phanerochaete chrysosporium* plays a central role in the initial degradation of the complex aromatic polymer lignin [1-4]. Lignin peroxidase catalyses reactions not normally associated with other peroxidases, that is, oxidation of non-phenolic aromatic substrates. The plethora of products formed from its substrates have been explained as the consequence of multiple, substituent-dependent, reactions taking place after the initial formation of aromatic cation radicals [5-10]. The same products have been found in experiments with non-enzymatic one-electron oxidants [11,12].

The methoxybenzenes, of which there are 12 congeners, were particularly useful as substrates in elucidating the cation-radical mechanism of lignin peroxidase [7]. In the presence of H_2O_2 , the enzyme oxidizes 10 of the 12 congeners, forming from several of them relatively stable cation radicals [7]. From the kinetics of radical formation, and results of product analyses, we concluded that oxidation of 1,4-dimethoxybenzene by lignin peroxidase results in demethoxylation to give *p*-benzoquinone and methanol as products. In studies with $H_2^{18}O$ we have shown that the quinone oxygen atoms come from water (P. J. Kersten, unpublished work).

The half-wave potentials (E_1) of the methoxybenzenes have been determined: these range from 0.81 V to 1.76 V versus a saturated calomel electrode [13]. This allows relationships to be examined between the redox potentials of the substrates and the

ability of enzymes to oxidize them by one electron. Hammel *et al.* [5] reported that lignin peroxidase oxidizes polycyclic aromatics with ionizing potentials of approx. 7.5 eV or lower. Cavalieri & Rogan [14] reported that HRP will oxidize only those polycyclics with ionization potentials below 7.35 eV.

In the present study we compared lignin peroxidase from *P. chrysosporium*, horseradish peroxidase (HRP) and laccase from the ligninolytic fungus *Trametes* (syn. *Coriolus* or *Polyporus*) *versicolor* in their oxidation of methoxybenzenes. HRP was included because it has a catalytic cycle very similar to that of lignin peroxidase [15,16], and the redox potentials of HRP compound I/compound II and compound II/ferric couples have been well characterized [17]. In contrast with the peroxidases, laccase is a Blue Copper oxidase that catalyses the four-electron reduction of O_2 to H_2O during its oxidation of phenolics, aromatic amines, ascorbate and metal cyanides [18]. Although some substrates are potential two-electron donors, the Blue oxidases are reduced in one-electron steps. This is due to the reduction mechanism in which Type 1 copper is the primary electron acceptor. The substrate radicals thus formed take part in further non-enzymic reactions [19,20]. The laccase from *T. versicolor* has Type 1 and Type 3 copper sites of particularly high redox potential in comparison with other Blue Copper proteins [21,22]. Furthermore, evidence has been presented recently that laccase, like lignin peroxidase, plays a role in lignin degradation by fungi [23], which makes additional information on its mode of action of interest to that area of research. A preliminary account of our results with lignin peroxidase and HRP has been presented [24].

Abbreviations used: HRP, horseradish peroxidase; E_1 , half-wave potential.

|| Present address: University of Wisconsin Biotechnology Center, Madison, WI 53705, U.S.A.

¶ To whom correspondence and reprint requests should be sent, at the following address: Forest Products Laboratory, Forest Service, U.S. Department of Agriculture, One Gifford Pinchot Drive, Madison, WI 53705-2398, U.S.A.

MATERIALS AND METHODS

Enzymes

Lignin peroxidase isoenzyme H8 was purified [25] from the extracellular broth of cultures of *Phanerochaete chrysosporium* [26]. It had a specific activity of 22 units \cdot mg⁻¹ on the basis of the oxidation of veratryl alcohol at pH 3 [27]. Protein determination of purified H8 was based on an ϵ_{400} of 168 mm⁻¹ \cdot cm⁻¹ [15]. HRP (Type VI) was purchased from Sigma Chemical Co. (St. Louis, MO, U.S.A.) and used without further purification. The two chromatographic forms, A and B, of laccase from *Trametes versicolor* were prepared according to previously published methods [28,29]. Both laccase forms were carefully characterized by optical-absorption, c.d. and e.s.r. spectroscopy. These studies excluded the possibility of any contaminating peroxidases [22].

Chemicals

Methoxybenzenes and 2,5-dimethoxy-*p*-benzoquinone were synthesized or purchased as previously described [7]. 4,5-Dimethoxy-*o*-benzoquinone was synthesized from catechol by the method of Wanzlick & Jahnke [30] [m.p. 224–227 °C (literature value 225–227 °C)].

U.v.-visible studies

To test whether a methoxybenzene congener was a substrate, as determined by u.v.-visible spectral analysis, enzyme reactions were run in 25 mM-sodium tartrate, pH 3.0, with a congener concentration of 2×10^{-4} M and 10 μ g of enzyme per ml of reaction mixture. With lignin peroxidase and HRP, the reactions were initiated with 4×10^{-4} M-H₂O₂. To monitor the decay of the 450-nm chromophore produced during the oxidation of 1,2,4,5-tetramethoxybenzene, 1×10^{-4} M-substrate and 2×10^{-4} M-H₂O₂ were used. For the determination of enzyme kinetic parameters with 1,2,4,5-tetramethoxybenzene at 450 nm ($\epsilon_{450} = 9800$ M⁻¹ \cdot cm⁻¹), enzyme levels were used so that the initial velocities were much greater than the rate of cation-radical decay. For the kinetic studies with pentamethoxybenzene, oxidation was monitored at 300 nm; a $\Delta\epsilon_{300}$ of 10800 M⁻¹ \cdot cm⁻¹ was determined for this oxidation. Because the reactions of the peroxidases are bisubstrate, saturating H₂O₂ (0.5 mM) was used to determine the kinetic parameters for the methoxybenzenes. Reactions with laccase were air-saturated.

E.s.r. measurements

E.s.r. measurements were performed as previously described [7], with reaction conditions the same as those described above for 'U.v.-visible studies'.

Product identification and quantification

2,5-Dimethoxy-*p*-benzoquinone and 4,5-dimethoxy-*o*-benzoquinone produced from the oxidation of 1,2,4,5-tetramethoxybenzene were identified and quantified with a Gilson (Middleton, WI, U.S.A.) System 41 Gradient Analytical h.p.l.c. apparatus equipped with a Vydac (Hesperia, CA, U.S.A.) reverse-phase Ultrasphere ODS column. A methanol/water (1:4, v/v) solvent was used at 1 ml/min and the absorbance monitored at 282 nm. Reaction conditions were as described above under 'U.v.-visible studies', with a reaction time of 2 h. Authentic samples were used for standardization. Methanol was also quantified by using alcohol oxidase from the yeast *Pichia pastoris* (Sigma) and a YSI model 53 oxygen monitor (Yellow Springs Instrument Co., Yellow Springs, OH, U.S.A.) fitted with a Gilson single-port 1.5 ml reaction chamber. The pH of the HRP-catalysed reaction mixture was adjusted to ~7.2 by the addition of 2 M-sodium phosphate, pH 7.8, after a 1 h incubation, and the

dissolved-O₂-levels monitored upon addition of 5 μ g of catalase (Worthington Biochemical Corp., Freehold, NJ, U.S.A.) and then 6 units of alcohol oxidase to 1.5 ml of the solution in the O₂-monitor chamber. Controls included the complete reaction mixture minus added H₂O₂.

¹⁸O-labelling studies

To test for ¹⁸O incorporation from H₂¹⁸O, reaction mixtures contained 15 μ g of HRP, 6×10^{-4} M-1,2,4,5-tetramethoxybenzene, and 1.2×10^{-3} M-H₂O₂ in 0.5 ml of tartrate buffer, pH 3.0, with either water or 10.6% excess ¹⁸O in water by dilution of 20% H₂¹⁸O (Cambridge Isotope Laboratories, Woburn, MA, U.S.A.). A control reaction was also run in which 2,5-dimethoxy-*p*-benzoquinone was substituted for tetramethoxybenzene to test for non-enzymic exchange of the quinone oxygen atoms with water. After 12 min the reactions were stopped and the quinones reduced with Na₂S₂O₄. The reduced products were extracted into dichloromethane, the extracts dried over anhydrous Na₂SO₄, the solvent evaporated, and the residues acetylated with acetic anhydride/pyridine (1:1, v/v). Excess reagents were removed under vacuum, the residue redissolved in dichloromethane, and analysed by g.c./m.s. using a Finnigan MAT (San Jose, CA, U.S.A.) 4510 instrument with a 60 m, 0.2 μ m-film-thickness (biscyanopropyl polysiloxane) SP-2340 fused-silica capillary column (Supelco, Bellefonte, PA, U.S.A.). The *m/z* values (relative intensity) at 70 eV for the enzymic reactions were as follows: in H₂O: 255 [*M*⁺ (the molecular ion)+1, 0.15%], 254 (*M*⁺, 9.1%), 212 (*M*⁺ minus H₂C=C=O, 11%), 171 (6.7%), 170 (*M*⁺ minus 2H₂C=C=O, 100%), 169 (2.0%), 156 (1.4%), 155 (44%); and in 10.6% H₂¹⁸O: 256 (3.6%), 255 (1.2%), 254 (12%), 214 (3.5%), 213 (1.2%), 212 (13%), 174 (1.6%), 173 (2.6%), 172 (28%), 171 (10%), 170 (100%), 157 (12%), 156 (3.3%), 155 (39%). The ¹⁸O content of the 2,5-dimethoxy-*p*-benzoquinone product was determined by comparing the ion counts at *m/z* 170, 172, and 174.

RESULTS

U.v.-visible spectral studies

To investigate the possibility that methoxybenzenes are substrates for HRP and laccase, we looked for u.v.-visible spectral changes upon incubation of the enzymes with the substrates as described in the Materials and methods section. Of the 12 methoxybenzenes, only the 1,2,4,5-congener was oxidized by both enzymes (Table 1). The three congeners closest to 1,2,4,5-tetramethoxybenzene in half-wave potential also gave spectral changes with HRP, but these clearly occurred more slowly than the rapid oxidation of the 1,2,4,5-congener. In comparison, lignin peroxidase was able to oxidize 10 of the 12 methoxybenzene congeners under similar conditions.

The spectral changes with HRP/H₂O₂ and 1,2,4,5-tetramethoxybenzene are shown in Fig. 1; similar results were obtained with laccase and lignin peroxidase. Interestingly, this oxidation resulted in biphasic spectral changes, with the rapid formation of an intermediate having a well-defined peak at 450 nm, which then slowly disappeared with apparent first-order kinetics. Plots of log absorbance versus time indicated a half-life for the 450 nm chromophore of 9.7 min, and extrapolation to zero time indicated an absorption coefficient (ϵ_{450}) of 9800 M⁻¹ \cdot cm⁻¹. Isosbestic points at 254 nm and 296 nm indicate that this intermediate is converted to final products without detectable accumulation of additional intermediates.

E.s.r. characterizations

The cation radical of 1,2,4,5-tetramethoxybenzene was identified by e.s.r. after oxidation by HRP and by laccase. The

Table 1. Oxidation of methoxybenzenes by lignin peroxidase, HRP and laccase

Compound	E_1 (V)*	Oxidation† by:		
		H8/H ₂ O ₂	HRP/H ₂ O ₂	Laccase/O ₂
Methoxybenzene	1.76	-	-	-
1,3-Dimethoxybenzene	ND	-	-	-
1,3,5-Trimethoxybenzene	1.49	±	-	-
1,2-Dimethoxybenzene	1.45	+	-	-
1,2,3-Trimethoxybenzene	1.42	+	-	-
1,4-Dimethoxybenzene	1.34	+‡	-	-
1,2,3,4-Tetramethoxybenzene	1.25	+‡	-	-
Hexamethoxybenzene	1.24	+‡	-	-
1,2,4-Trimethoxybenzene	1.12	+	+	-
1,2,3,5-Tetramethoxybenzene	1.09	+	+	-
Pentamethoxybenzene	1.07	+	+	-
1,2,4,5-Tetramethoxybenzene	0.81	+‡	+‡	+‡

* E_1 values are literature values [13] obtained by polarography with respect to a saturated calomel electrode; ND, not determined.

† A '+ ' indicates that u.v.-visible spectral changes were observed with enzyme and H₂O₂, whereas '-' indicates that they were not. A '±' indicates that spectral changes were observed, but the enzyme was quickly inactivated.

‡ Cation radical detected directly by e.s.r.

hyperfine coupling parameters were as reported previously for the oxidation by lignin peroxidase [7] and as reported for the cation radical produced chemically [31]. The spectrum generated by oxidation with HRP/H₂O₂ is shown in Fig. 2 along with an inset showing the first-order decay of the cation radical; it has

the same half-life as the 450 nm chromophore described above, indicating that the cation radical is the intermediate species detected in the u.v.-visible studies.

The oxidation of 1,3,5-trimethoxybenzene by lignin peroxidase H8 was unusual, not only because the enzyme was rapidly inactivated during the oxidation, but also because it was the only oxidation in the series for which a radical species could easily be detected by e.s.r. spectroscopy using 5,5-dimethyl-1-pyrroline *N*-oxide as spin-trap. The e.s.r. spectra were consistent with trapping of a carbon-centred radical in the absence of O₂ and an O₂-centred radical in air (results not shown). It is conceivable that inactivation of lignin peroxidase in air is linked to the production of peroxy radicals, as has been suggested for horseradish peroxidase [32].

Product analyses

On the basis of our previous results with lignin peroxidase and 1,4-dimethoxybenzene, an expected product from decay of the




Fig. 1. U.v.-visible spectral changes in 1,2,4,5-tetramethoxybenzene on incubation with HRP and H₂O₂.

Upon initiation of the reaction with H₂O₂, the spectrum of the starting material (290 nm λ_{max} , lowest spectrum) was quickly replaced with a new spectrum of λ_{max} 290 nm (shoulder: 305 nm) and λ_{max} 450 nm. The absorbance at 450 nm and at 305 nm then slowly decreased to give a new peak at 282 nm. The cycle time was 3.12 min. The reaction was run in 25 mM-sodium tartrate, pH 3, with 8×10^{-5} M-1,2,4,5-tetramethoxybenzene, 10 μ g of HRP/ml of reaction mixture and 3.3×10^{-4} M-H₂O₂.




Fig. 2. E.s.r. signal of the cation radical of 1,2,4,5-tetramethoxybenzene, obtained by oxidation with HRP and H₂O₂.

A similar spectrum was obtained with laccase with magnetic parameters as earlier determined with lignin peroxidase [7]. The inset shows the first-order decay of the signal, indicating a half life of 9.97 min. Intensity is given in arbitrary units.




Fig. 3. Mass-spectral analyses of ^{18}O -labelled products

2,5-Dimethoxy-*p*-benzoquinone from the oxidation of 1,2,4,5-tetramethoxybenzene by HRP/ H_2O_2 was reduced and acetylated before analysis. Reactions were run in either natural-abundance H_2O (a) or enriched in ^{18}O (b). A control with authentic 2,5-dimethoxy-*p*-benzoquinone incubated in an H_2^{18}O -reaction mixture showed no enrichment for ^{18}O .




Fig. 4. pH optimum for the oxidation of 1,2,4,5-tetramethoxybenzene by HRP and laccase A

Reaction mixtures contained 7.5 mM-1,2,4,5-tetramethoxybenzene and either 50 ng of HRP (■) or 0.25 μg of laccase A (▲) in 1 ml of 50 mM-tartrate/citrate buffer. HRP-catalysed reactions were initiated with 0.5 mM- H_2O_2 . Initial velocities were determined at 450 nm (see the Materials and methods section).




Fig. 5. pH optimum for the oxidation of pentamethoxybenzene by lignin peroxidase and HRP

Reaction mixtures contained 2 mM-pentamethoxybenzene, 0.5 mM- H_2O_2 and either 1.68 μg of purified H8 lignin peroxidase (■) or 10 μg HRP (▲) in 1 ml of 50 mM-tartrate buffer. In both cases, activity was less stable at pH values below 3.

Table 2. Steady-state kinetic constants for the oxidation of 1,2,4,5-tetramethoxybenzene and pentamethoxybenzene

Apparent kinetic parameters were determined spectroscopically as described in the Materials and methods section for reactions in 50 mM-sodium tartrate, pH 3.0. Oxidations of 1,2,4,5-tetramethoxybenzene were monitored by measuring the production of the cation radical; rates are therefore corrected for a two-electron oxidation.

Enzyme	1,2,4,5-Tetramethoxybenzene			Pentamethoxybenzene		
	K_m (mM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1}\cdot\text{s}^{-1}$)	K_m (mM)	k_{cat} (s^{-1})	k_{cat}/K_m ($\text{M}^{-1}\cdot\text{s}^{-1}$)
H8	0.09	11	1.2×10^5	0.12	9.9	8.2×10^4
HRP	1.2	625	5.2×10^5	> 7.5*	> 2.3*	307*
Laccase A	2.2	120	5.5×10^4	—†	—	—
Laccase B	1.5	124	8.3×10^4	—	—	—

* Non-saturating conditions were observed and therefore the second-order rate constant was determined by dividing the observed rate by enzyme and substrate concentrations.

† No oxidation detected.

cation radical of 1,2,4,5-tetramethoxybenzene would be 2,5-dimethoxy-*p*-benzoquinone [7]. In accord with this expectation, this quinone, as well as 4,5-dimethoxy-*o*-benzoquinone, was identified and quantified by h.p.l.c. The two products were formed in a ratio of approx. 4 (*p*-isomer):1 (*o*-isomer) and

accounted for all the ring-derived product within experimental error. Virtually the same qualitative and quantitative results were obtained with lignin peroxidase, HRP and laccase. Furthermore, an experiment with HRP and H_2^{18}O showed that the quinone oxygen atoms are derived from water (see Fig. 3). We also

determined that 2 mol (1.9 ± 0.2) of methanol are formed from each 1 mol of substrate oxidized.

pH optima

As a prelude to steady-state kinetic studies, the pH optima for the oxidation of 1,2,4,5-tetramethoxybenzene by HRP and laccase (Fig. 4) and of pentamethoxybenzene by HRP and lignin peroxidase (Fig. 5) were examined. Surprisingly, in all cases a low pH optimum was observed for the initial velocities. A similar pH-optimum profile was obtained for laccase A assayed with a Clark-type oxygen electrode (results not shown).

Steady-state kinetics

The data in Table 1 indicate substantial thermodynamic differences among the three enzymes. It was of interest therefore to compare kinetic parameters with substrates common to the enzymes, i.e. with methoxybenzenes of the lowest E_i values. On the basis the pH-optima results above, and on previous investigations of the kinetic parameters of lignin peroxidase [15,33], the steady-state kinetics for the oxidation of 1,2,4,5-tetramethoxybenzene and pentamethoxybenzene were determined at pH 3 (Table 2). A saturating level of H_2O_2 (0.5 mM) was used with the peroxidases.

DISCUSSION


Our results show that 1,2,4,5-tetramethoxybenzene is oxidized by HRP and laccase in the same way as it is by lignin peroxidase, namely via formation of the aryl cation radical. With all three enzymes, the cation radical degrades to the same products, indicating that the enzymes all have the same effect. Results with $H_2^{18}O$ and the HRP reaction showed that the quinone oxygen atoms are derived from water, as they are in the lignin peroxidase-catalysed oxidation of the 1,4-congener (P. J. Kersten, unpublished work). On the basis of these results,

we propose the sequence shown in Scheme 1 for the oxidation of 1,2,4,5-tetramethoxybenzene by the three enzymes.

There has been very little work to date on *O*-demethylation reactions catalysed by typical (non-ligninolytic) peroxidases such as HRP. The *O*-demethylation of 9-methoxyellipticine derivatives by HRP/ H_2O_2 [34] resembles that of the methoxybenzenes [7; the present study] in that the methyl group is lost as methanol and the quinone oxygen atom of the quinone-imine products is derived from water. In the 9-methoxyellipticine case, however, the authors [34] proposed that the addition of water to intermediates occurs after two-electron oxidation (and loss of a proton) without the involvement of cation-radical intermediates.

With the homologous series of methoxybenzenes, oxidizability by the three enzymes correlated with E_i of the compounds and not with the methoxyl substitution pattern (see Table 1). Because the E_i values in Table 1 were obtained in acetonitrile with respect to a saturated calomel electrode [13], conclusions about the magnitude of these potentials in a different environment should be made with caution. We presume, however, that with the homologous series of substrates, solvent effects for the various congeners are similar and that comparisons within the series remain valid.

Our results with the methoxybenzenes indicate that all three enzymes oxidize their aromatic substrates by removing one electron from the aromatic nuclei. Apparently this is the first report of oxidation by laccase of aromatic nuclei bearing no hydroxy or amino group, although it is well known that lignin peroxidase oxidizes such substrates [1-5], and HRP oxidation of polycyclic aromatics has been reported [14]. Electron-donating groups such as hydroxy groups and amines, however, would be expected to facilitate oxidation by lowering the redox potential of the aromatic nuclei, which our results indicate is a critical parameter. In accord, all three enzymes oxidize a broad range of aromatic amines and phenols. This broad substrate specificity of



Scheme 1. Hypothetical scheme for the oxidation of 1,2,4,5-tetramethoxybenzene by lignin peroxidase/ H_2O_2 , HRP/ H_2O_2 or laccase/ O_2 .

the three enzymes suggests that they do not have specific binding sites.

The results also indicate that lignin peroxidase is a stronger oxidant than HRP, which in turn is stronger than laccase. The conclusion that oxidized HRP is a better oxidant than laccase is consistent with the published potentials for these enzymes. Using equilibria data from reactions coupled with the K_2IrCl_6 - K_3IrCl_6 system, Hayashi & Yamasaki [17] determined redox potentials of ~950 mV for HRP compound I/compound II and compound II/ferric couples at slightly acidic pH values. From potentiometric studies using metal cyanides as electron mediators, Reinhammar [21] determined redox potentials of 785 and 782 mV for the Type 1 and Type 3 copper sites, respectively, of the laccase from *T. versicolor* at pH 5.5. The data of Table 1 would suggest higher redox values for these enzymes, probably owing to the lower pH and non-equilibrium conditions employed in our experiments. An oxidation potential more than 250 mV above that of oxidized HRP is suggested for oxidized lignin peroxidase at pH 3.

The kinetics for oxidation of the congeners of lowest redox potential demonstrate further differences in the enzymes. Notably, the K_m values for 1,2,4,5-tetramethoxybenzene and pentamethoxybenzene with lignin peroxidase H8 are at least an order of magnitude lower than those observed with HRP and laccase. These K_m values observed with H8, however, are in reasonable agreement with results obtained for veratryl alcohol, 1,4-dimethoxybenzene and the lignin model compound 1-(3,4-dimethoxyphenyl)-2-(2-methoxyphenoxy)-propane-1,3-diol, which represents the β -O-4 substructure of lignin [35]. In each case, a K_m value between 50 μ M and 150 μ M was determined. Likewise, the k_{cat} values observed here are of the same order of magnitude.

1,2,4,5-Tetramethoxybenzene appears to be a better substrate for HRP than it is for lignin peroxidase. Despite the relatively high K_m for this substrate with HRP, the k_{cat} value observed is about 60 times greater than that with H8, indicating a second-order rate constant (k_{cat}/K_m) that is about four times greater for HRP than for H8. Likewise, the laccases have ten-fold higher k_{cat} values than lignin peroxidase with 1,2,4,5-tetramethoxybenzene; the k_{cat}/K_m values indicate that lignin peroxidase is only a slightly more efficient catalyst than laccase in this case. However, with pentamethoxybenzene a very different relationship is observed; lignin peroxidase is a much more efficient catalyst than HRP, and no activity is observed with the laccases.

The order of k_{cat} values for the three enzymes for the oxidation of the 1,2,4,5-congener is not the same as expected on the basis of the relative oxidation potentials of the enzymes. That is, k_{cat} (and k_{cat}/K_m as well) clearly depends on factors in addition to the oxidation potential of the enzymes. Such factors would be expected to include distance between interacting redox centres, orientation of substrate in or on the enzyme, and steric and electronic influences of neighbouring groups of the enzymes and substrates during oxidation [36].

Interestingly, the oxidation of 1,2,4,5-tetramethoxybenzene by all three enzymes was favoured at low pH. The low pH optima with the non-phenolic substrates suggest that oxidation is controlled by substrate-specific factors (e.g. stabilization of intermediates) that are favoured at low pH. The transient-state kinetics of lignin peroxidase compound I and compound II oxidation of veratryl alcohol likewise show a strong pH-dependence with faster oxidations at lower pH values [37].

We acknowledge the help of Mr. Mike Mozuch and Dr. Roger Peterson of the Forest Products Laboratory in performing h.p.l.c. and

m.s. analyses respectively. This work was supported in part by National Institutes of Health grants GM-29035 and RR-018008.

REFERENCES

1. Buswell, J. A. & Odier, E. (1987) *CRC Crit. Rev. Biotechnol.* **6**, 1-60
2. Higuchi, T. (1986) *Wood Res. (Kyoto Univ.)* **73**, 58-81
3. Kirk, T. K. & Farrell, R. L. (1987) *Annu. Rev. Microbiol.* **41**, 465-505
4. Tien, M. (1987) *CRC Crit. Rev. Microbiol.* **15**, 141-168
5. Hammel, K. E., Kalyanaraman, B. & Kirk, T. K. (1986) *J. Biol. Chem.* **261**, 16948-16952
6. Hammel, K. E., Tien, M., Kalyanaraman, B. & Kirk, T. K. (1985) *J. Biol. Chem.* **260**, 8348-8353
7. Kersten, P. J., Tien, M., Kalyanaraman, B. & Kirk, T. K. (1985) *J. Biol. Chem.* **260**, 2609-2612
8. Kirk, T. K., Tien, M., Kersten, P. J., Mozuch, M. D. & Kalyanaraman, B. (1986) *Biochem. J.* **236**, 279-287
9. Renganathan, V., Miki, K. & Gold, M. H. (1985) *Arch. Biochem. Biophys.* **246**, 155-161
10. Shoemaker, H. E., Harvey, P. J., Bowen, R. M. & Palmer, J. M. (1985) *FEBS Lett.* **183**, 7-12
11. Shimada, M., Habe, T., Higuchi, T., Okamoto, T. & Panijpan, B. (1987) *Holzforchung* **41**, 277-285
12. Huynh, V.-B. (1986) *Biochem. Biophys. Res. Commun.* **139**, 1104-1110
13. Zweig, A., Hodgson, W. G. & Jura, W. H. (1964) *J. Am. Chem. Soc.* **86**, 4124-4129
14. Cavalieri, E. & Rogan, E. (1985) *Environ. Health Perspect.* **64**, 69-84
15. Tien, M., Kirk, T. K., Bull, C. & Fee, J. A. (1986) *J. Biol. Chem.* **261**, 1687-1693
16. Renganathan, V. & Gold, M. H. (1986) *Biochemistry* **25**, 1626-1631
17. Hayashi, Y. & Yamasaki, I. (1979) *J. Biol. Chem.* **254**, 9101-9106
18. Malmström, B. G., Andréasson, L.-E. & Reinhammar, B. (1975) in *The Enzymes* (Boyer P. D., ed.), vol. 12, pp. 507-579, Academic Press, New York
19. Andréasson, L.-E. & Reinhammar, B. (1976) *Biochim. Biophys. Acta* **445**, 579-597
20. Andréasson, L.-E. & Reinhammar, B. (1979) *Biochim. Biophys. Acta* **568**, 145-156
21. Reinhammar, B. (1972) *Biochim. Biophys. Acta* **275**, 245-259
22. Reinhammar, B. (1984) in *Copper Proteins and Copper Enzymes* (Lontie, R., ed.), pp. 1-35, CRC Press, Boca Raton, FL
23. Kawai, S., Umezawa, T. & Higuchi, T. (1988) *Arch. Biochem. Biophys.* **262**, 99-110
24. Kersten, P. J., Kalyanaraman, B., Hammel, K. E. & Kirk, T. K. (1987) in *Lignin Enzymic Microbial Degradation* (International Symposium Odier, E., ed.), pp. 107-122, Institut National de la Recherche Agronomique, Paris
25. Kirk, T. K., Croan, S., Tien, M., Murtagh, K. E. & Farrell, R. L. (1986) *Enzyme Microb. Technol.* **8**, 27-32
26. Jäger, A., Croan, S. & Kirk, T. K. (1985) *Appl. Environ. Microbiol.* **50**, 1274-1278
27. Tien, M. & Kirk, T. K. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**, 2280-2284
28. Fähræus, G. & Reinhammar, B. (1967) *Acta Chem. Scand.* **21**, 2367-2378
29. Nestor, L., Reinhammar, B. & Spiro, T. G. (1986) *Biochim. Biophys. Acta* **869**, 286-292
30. Wanzlick, H. W. & Jahnke, U. (1968) *Chem. Ber.* **101**, 3744-3752
31. Ishizu, K., Watanabe, K. & Ohya-Nishiguchi, H. (1980) in *Landolt-Börnstein* (Fischer, H. & Hellwege, K.-H., eds.), group 2, vol. 9, pp. 124-147, Springer-Verlag, New York
32. Ma, X. & Rokita, S. E. (1988) *Biochem. Biophys. Res. Commun.* **157**, 160-165
33. Andrawis, A., Johnson, K. A. & Tien, M. (1988) *J. Biol. Chem.* **263**, 1195-1198
34. Meunier, G. & Meunier, B. (1985) *J. Biol. Chem.* **260**, 10576-10582
35. Farrell, R. L., Murtagh, K. E., Tien, M., Mozuch, M. D. & Kirk, T. K. (1989) *Enzyme Microb. Technol.* **11**, 322-328
36. Marcus, R. A. & Sutin, N. (1985) *Biochim. Biophys. Acta* **881**, 265-322
37. Marquez, L., Wariishi, H., Dunford, H. B. & Gold, M. H. (1988) *J. Biol. Chem.* **263**, 10549-10552