

REVIEW ARTICLE

Sunscreens: are they beneficial for health? An overview of endocrine disrupting properties of UV-filters

M. Krause,*† A. Klit,† M. Blomberg Jensen,* T. Søbørg,* H. Frederiksen,* M. Schlumpf,‡ W. Lichtensteiger,‡ N. E. Skakkebaek* and K. T. Drzewiecki†

*Department of Growth and Reproduction, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark, †Department of Plastic surgery, Breast surgery and Burns, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark, and ‡GREEN Tox and Institute of Anatomy, University of Zurich, Winterthurerstrasse 190, Zurich, Switzerland

Summary

Keywords:

benzophenone-3, endocrine disruptors, homosalate, malignant melanoma, OD-PABA, PABA, sunscreens, UV-filters, 3-benzylidene camphor, 3-(4-methyl-benzylidene) camphor, 2-ethylhexyl 4-methoxy cinnamate

Correspondence:

N.E. Skakkebaek, Department of Growth and Reproduction, University of Copenhagen, Rigshospitalet, Copenhagen, Denmark.
E-mail: nes@rh.dk

Received 5 January 2012; revised 13 March 2012; accepted 13 March 2012

doi:10.1111/j.1365-2605.2012.01280.x

Today, topical application of sunscreens, containing ultraviolet-filters (UV-filters), is preferred protection against adverse effects of ultraviolet radiation. Evidently, use of sunscreens is effective in prevention of sunburns in various models. However, evidence for their protective effects against melanoma skin cancer is less conclusive. Three important observations prompted us to review the animal data and human studies on possible side effects of selected chemical UV-filters in cosmetics. (1) the utilization of sunscreens with UV-filters is increasing worldwide; (2) the incidence of the malignant disorder for which sunscreens should protect, malignant melanoma, is rapidly increasing and (3) an increasing number of experimental studies indicating that several UV-filters might have endocrine disruptive effects. The selected UV-filters we review in this article are benzophenone-3 (BP-3), 3-benzylidene camphor (3-BC), 3-(4-methyl-benzylidene) camphor (4-MBC), 2-ethylhexyl 4-methoxy cinnamate (OMC), Homosalate (HMS), 2-ethylhexyl 4-dimethylaminobenzoate (OD-PABA) and 4-aminobenzoic acid (PABA). The potential adverse effects induced by UV-filters in experimental animals include reproductive/developmental toxicity and disturbance of hypothalamic–pituitary–thyroid axis (HPT). Few human studies have investigated potential side effects of UV-filters, although human exposure is high as UV-filters in sunscreens are rapidly absorbed from the skin. One of the UV-filters, BP-3, has been found in 96% of urine samples in the US and several UV-filters in 85% of Swiss breast milk samples. It seems pertinent to evaluate whether exposure to UV-filters contribute to possible adverse effects on the developing organs of fetuses and children.

Abbreviations

3-BC, 3-benzylidene camphor; 4-MBC, 3-(4-methyl-benzylidene)-camphor; ↑, Increased; ↓, Decreased; AhR, aryl hydrocarbon receptor; AR, androgen receptor; BP-3, benzophenone-3; bw, body weight; C3, complement protein 3; Dio1, 5′deiodinase type I; EDC, endocrine disrupting chemicals; ER, oestrogen receptor; ERR1, oestrogen receptor related receptor 1; F0, parent rats; F1, 1. generation of rat offspring; FDA, Food and Drug Administration; FRTL-5, normal, non-transformed rat thyrocytes; FT3, free triiodothyronine; FT4, free thyroxine; hAR, human androgen receptor; HepG2, human hepatocarcinoma cell line; hER, human oestrogen receptor; HMS, homosalat; hPR, human progesterone receptor; HPT, hypothalamic–pituitary–thyroid axis; hrTPO, human recombinant thyroid peroxidase; IGF-I, insulin-like growth factor-I; LDH, lactate dehydrogenase; LOAEL, lowest observed adverse effect levels; M, male; MCF7, human breast cancer cells; ME, malic enzyme; MM, malignant melanoma; MPO, medial preoptic area; N-Cor, nuclear receptor co-repressor; NHANES, National Health and Nutrition Examination Survey; NOAEL, no observed adverse effect levels; OCT, 2-cyano-3,3-diphenyl acrylic acid; OD-PABA, 2-ethylhexyl 4-dimethylaminobenzoate; OMC, 2-ethylhexyl-4-methoxy cinnamate; P, plasma; ORG2058, PR agonist; PABA, 4-aminobenzoic acid; PND, post natal day; PND1, day of birth; PR, progesterone receptor; rtER, rainbow trout oestrogen receptor; SRC-1, steroid receptor coactivator-1; T3, total triiodothyronine; T4, total thyroxine; TBG, thyroxine-binding globulin; TPO, thyroid peroxidase; TR, thyroid hormone receptor; TRH, thyrotropin releasing hormone; TSH, thyroid-stimulating hormone; U2-OS, cells, human osteosarcoma cells; UV-A, ultraviolet radiation with wavelength A, 320–400 nm; UV-B, ultraviolet radiation with wavelength B, 290–320 nm; VMH, ventromedial hypothalamic nucleus- plays important role in sexual behaviour and receptivity of female rats (191); VTG, vitellogenin (oestrogen-responsive gene products in fish).

Introduction

The first commercial sunscreen was developed in the 1930s to abrogate ultraviolet-B waveband (UV-B), and thus prevent sunburn (Rebut, 1990). In 1970, sunscreens were developed further to protect against both ultraviolet-A waveband (UV-A) and UV-B (Deep, 2010), because of their suggested causal role in the development of skin cancer, in particular malignant melanoma (MM) (Wang *et al.*, 2001; Gandini *et al.*, 2005). Today it is still questionable whether this aim has been achieved. There is no doubt that sunscreens protect against sunburn, solar keratosis, and non-melanoma skin cancer (Thompson *et al.*, 1993; Green *et al.*, 1999; Dupuy *et al.*, 2005). However, the only randomized trial examining the risk of MM after regular sunscreen use, found borderline statistical significance for a reduced incidence of new primary melanoma (Green *et al.*, 2011). In addition, despite use of sunscreens with




Figure 1 Age standardized MM incidence in USA, AU, NZ, UK and DK (Local cancer statistics). DK: Denmark: 1975–2000 (IARC); 2001–2010: Danish Cancer registers (Sundhedsstyrelsen, 2012). UK: United Kingdom: (Cancer Research UK). USA: (SEER). AUS: Australia: (Australian Institute of Health and Welfare [AIHW]).

UV-filters over decades (Figure S1), the incidence of MM is still increasing rapidly (Fig. 1) (Handel & Ramagopalan, 2010). Furthermore, an increasing number of experimental animal and in vitro studies indicated that some UV-filters might have adverse effects as endocrine disrupters.

In light of the high incidence of MM and considering the facts that the protective effect of sunscreens against MM has not been fully proven, we have reviewed the literature on possible endocrine disrupting effects of the most common chemical UV-filters used in cosmetics. Those are benzophenone-3 (BP-3), 3-benzylidene camphor (3-BC), 3-(4-methyl-benzylidene) camphor (4-MBC), 2-ethylhexyl 4-methoxy cinnamate (OMC), homosalate (HMS), 2-ethylhexyl 4-dimethylaminobenzoate (OD-PABA) and 4-aminobenzoic acid (PABA).

In vitro and in vivo adverse effects of UV-filters

A wide range of in vitro and in vivo studies have identified several UV-filters as endocrine disrupting chemicals (EDC) (European Commission) (Table S1). Table S1 lists results from in vivo and in vitro studies where possible adverse effects of the most common chemical UV-filters in cosmetics (Table 1) were examined. In the following the main results from Table S1 are summarized. A simplified version of Table S1 is presented in Table 2.

Influence on oestrogenic signalling

In vitro studies, investigating oestrogenic activity of UV-filters, varied in their design and endpoints, which may explain the diverging results. The majority of the in vitro studies reported that BP-3, 4-MBC, OMC, HMC and OD-PABA all exhibit oestrogenic activity (Schlumpf *et al.*, 2001b, 2004a; Schreurs *et al.*, 2002, 2005; Gomez *et al.*, 2005; Kunz *et al.*, 2006). However, not all of the UV-filters exhibiting in vitro oestrogenic activity were

Table 1 Most common UV-filters in cosmetics

	Generic name	Product name	Max concentration (%)	Spectrum of action	Approved
Chemical UV-filters	Benzophenone-3	BP-3	6 ^b –10 ^{a,c}	UV-A, UV-B	EU, US, AU
	2-Cyano-3,3-diphenyl acrylic acid	OCT	10	UV-B	EU, US, AU
	3-Benzylidene camphor	3-BC	2	UV-B	EU
	3-(4-Methyl-benzylidene) camphor	4-MBC	4	UV-B	EU, AU
	2-Ethylhexyl 4-methoxy cinnamate	OMC	7.5 ^b –10 ^{a,c}	UV-B	EU, US, AU
	Homosalate	HMS	10 ^a –15 ^{b,c}	UV-B	EU, US, AU
	2-Ethylhexyl 4-dimethylaminobenzoate	OD-PABA	8	UV-B	EU, US, AU
	4-Aminobenzoic acid	PABA	15 ^{b,c}	UV-B	US, AU
Physical UV-filter	Titanium dioxide		25	Physical	EU, US, AU
	Zinc oxide		25-no limit	Physical	US, AU

^aList of permitted UV-filters in the Council Directive of the European Committee.

^bList of permitted UV-filters in the US Food and Drug Administration monograph.

^cList of permitted UV-filters in the Australian regulatory guidelines for over-the-counter medicines (ARGOM), by the Therapeutic Goods Administration.

Table 2 In vitro and In vivo effects of the most common UV-filters, used in cosmetics (increased: ↑; decreased: ↓)

Endpoint	References
BP-3	
Oestrogen activity: In vitro	
Binding to hER α	Morohoshi <i>et al.</i> (2005)
↑ Transactivation: ER α , ER β	Gomez <i>et al.</i> (2005), Kunz <i>et al.</i> (2006), Kunz & Fent (2006b), Morohoshi <i>et al.</i> (2005) and Schreurs <i>et al.</i> (2002)
Activation: hER β > hER α	Schreurs <i>et al.</i> (2005)
No Antagonism of ER β transactivation	Schreurs <i>et al.</i> (2002)
Antagonistic action > agonistic action on hER α	Kunz & Fent (2006b)
↑ MCF-7 cell proliferation	Schlumpf <i>et al.</i> (2001b)
Oestrogen activity: acute in vivo models	
Uterotrophic effect in immature rats	Schlumpf <i>et al.</i> (2001b)
Uterus of adult oophorectomized rats: unchanged weight, ↓ ER β , unchanged ER α , ERR1 and AhR expression	Schlecht <i>et al.</i> (2004)
No effect on VTG induction in juvenile fathead minnows	Kunz <i>et al.</i> (2006)
Androgen activity: In vitro	
Antagonism of hAR transactivation. No agonistic action	Kunz & Fent (2006b), Ma <i>et al.</i> (2003) and Schreurs <i>et al.</i> (2005)
Progesterone activity: In vitro	
Antagonism of hPR transactivation	Schreurs <i>et al.</i> (2005)
Reproductive organs: long-term exposure of adult animals	
Oral exposure : ↓epididymal sperm density (rat, mice), ↑abnormal spermatozoa (mice). Dermal exposure (mice): ↓epididymal sperm density	French (1992)
Dermal exposure (rats): no effect	
Oral exposure (mice); ↑oestrous cycle length, Dermal exposure (rats): no effect	French (1992)
Reproductive organs: developmental toxicity	
No data available	
Thyroid axis	
In vitro: ↑ TR transcription	Schmutzler <i>et al.</i> (2007b)
↓ ERR1, unchanged ER α , ER β and AhR expression in thyroid of adult oophorectomized rats (5 day)	Schlecht <i>et al.</i> (2004)
Additional organ toxicities and general toxicity	
↓ Food consumption (2 weeks), ↓ body weight, ↑ liver weight (13 weeks)	French (1992)

Table 2 Continued

Endpoint	References
BP-2	
Thyroid axis	
In vitro: ↓hrTPO activity, unaltered iodide uptake	Schmutzler <i>et al.</i> (2007a)
adult oophorectomized rats (5 days): ↑TSH, ↓T4, altered Dio 1 activity (liver), unaltered TPO activity	
3-BC	
Oestrogen activity: In vitro	
Binding to hER β , no binding to hER α	Schlumpf <i>et al.</i> (2004a)
↑ Transactivation: ER α	Kunz <i>et al.</i> (2006) and Kunz & Fent (2006b)
Activation: hER β > hER α	Schreurs <i>et al.</i> (2005)
Antagonistic action > agonistic action on hER α	Kunz & Fent (2006b)
↑ MCF-7 cell proliferation	Schlumpf <i>et al.</i> (2001b)
Oestrogen activity: Acute in vivo models	
Uterotrophic effect in immature rats (3 days)	Schlumpf <i>et al.</i> (2001b)
↑ VTG induction in juvenile rainbow trouts and juvenile fathead minnows	Holbech <i>et al.</i> (2002) and Kunz <i>et al.</i> (2006)
Androgen activity: In vitro	
No antagonism of hAR transactivation	Ma <i>et al.</i> (2003)
Antagonism of hAR transactivation. No agonistic action	Holbech <i>et al.</i> (2002), Kunz & Fent (2006b) and Schreurs <i>et al.</i> (2005)
Progesterone activity: In vitro	
Antagonism of hPR transactivation	Schreurs <i>et al.</i> (2005)
Reproductive organs: Developmental toxicity	
Delayed male puberty in F1	Faass <i>et al.</i> (2009) and Schlumpf <i>et al.</i> (2008)
Oestrous cycle changes in F1	Schlumpf <i>et al.</i> (2008)
↓ Weight of uterus (high dose) and prostate (low dose) F1	Schlumpf <i>et al.</i> (2004b)
Altered gene expression in uterus and prostate in F1	
Central nervous system: Developmental toxicity	
Impaired female sexual behaviour in F1	Faass <i>et al.</i> (2009) and Schlumpf <i>et al.</i> (2008)
Additional organ toxicities and general toxicity	
Body weight in adult F1 (highest dose) (developmental study)	Schlumpf <i>et al.</i> (2004b)
↓ Length and weight, juvenile fathead minnows (14 days)	Kunz <i>et al.</i> (2006)
HMS	
Oestrogen activity: In vitro	
↑ Transactivation: hER α ,	Gomez <i>et al.</i> (2005)
↑ Transactivation: hER α > hER β	Schreurs <i>et al.</i> (2002, 2005)

Table 2 *Continued*

Endpoint	References
HMS	
No agonistic action: hER α , rER α	Kunz <i>et al.</i> (2006) and Kunz & Fent (2006b)
Antagonism of hER α transactivation, No antagonism at hER α or hER β	Kunz & Fent (2006b) Schreurs <i>et al.</i> (2002)
↑ MCF-7 cell proliferation	Schlumpf <i>et al.</i> (2001b)
Oestrogen activity: Acute in vivo models	
No uterotrophic effect in immature rats	Schlumpf <i>et al.</i> (2001b)
Androgen activity: In vitro	
Antagonism of hAR transactivation. No agonistic action	Ma <i>et al.</i> (2003) and Schreurs <i>et al.</i> (2005)
Agonistic and antagonistic action on hAR transactivation	Kunz & Fent (2006b)
Progesterone activity: In vitro	
Antagonism of hPR transactivation	Schreurs <i>et al.</i> (2005)
4-MBC	
Oestrogen activity: In vitro	
Binding to cytosolic ER	Tinwell <i>et al.</i> (2002)
Binding to hER β	and Schlumpf <i>et al.</i> (2004a)
No binding to hER α	Morohoshi <i>et al.</i> (2005)
↑ transactivation: hER α , hER β	Gomez <i>et al.</i> (2005) and Schreurs <i>et al.</i> (2002)
Activation hER α > hER β	Schreurs <i>et al.</i> (2005)
Activation hER α < hER β	Mueller <i>et al.</i> (2003)
No transactivation: hER α , rER α	Kunz <i>et al.</i> (2006), Kunz & Fent (2006b) and Morohoshi <i>et al.</i> (2005)
Antagonism of hER	Kunz & Fent (2006b) and Mueller <i>et al.</i> (2003)
No antagonism of hER	Morohoshi <i>et al.</i> (2005) and Schreurs <i>et al.</i> (2002)
↑ ER-mediated MCF-7 cell proliferation	Schlumpf <i>et al.</i> (2001b, 2004a) and Tinwell <i>et al.</i> (2002)
Oestrogen activity: Acute in vivo models	
Uterotrophic effect in immature rats	Schlumpf <i>et al.</i> (2001b) and Tinwell <i>et al.</i> (2002)
Uterotrophic effect in immature hairless Nu rats (dermal exposure)	Schlumpf <i>et al.</i> (2001b)
VTG induction in juvenile fathead minnows: no effect	Kunz <i>et al.</i> (2006)
No effect in transgenic juvenile zebra fish	Schreurs <i>et al.</i> (2002)
Androgen activity: In vitro	
Antagonism of hAR transactivation. No agonistic action	Kunz & Fent (2006b)
No antagonism of hAR transactivation	Ma <i>et al.</i> (2003) and Schreurs <i>et al.</i> (2005)

Table 2 *Continued*

Endpoint	References
4-MBC	
Progesterone activity: In vitro	
Antagonism of hPR transactivation	Schreurs <i>et al.</i> (2005)
Reproductive organs: Developmental toxicity	
Delayed puberty in males (prepubertal separation),	Durrer <i>et al.</i> (2005, 2007) and Hofkamp <i>et al.</i> (2008)
↑ prostate duct formation in F1 neonate,	
↓ prostate weight in adult F1,	
↑ testis weight in adult F1,	
↑ uterine weight	
Altered expression and sensitivity of oestrogen target genes and coactivators in prostate and uterus, no effect on onset of female puberty or oestrous cycle	Durrer <i>et al.</i> (2005, 2007) and Faass <i>et al.</i> (2009)
Reproductive organs: Long-term exposure of adult animals	
Uterus and vagina of adult oophorectomized rats	Seidlova-Wuttke <i>et al.</i> (2006)
↑ uterus weight, ↑ epithelial/ endometrial thickness, unchanged ER, PR and IGF-1 expression	
Central nervous system: Developmental toxicity	
Impaired female sexual behaviour in adult F1	Faass <i>et al.</i> (2009), Maerkel <i>et al.</i> (2005, 2007)
Altered expression and sensitivity of oestrogen target genes in sexually dimorphic brain regions	
Thyroid axis	
in vitro:	Schmutzler <i>et al.</i> (2007b)
↓ Iodide uptake	
Developmental study, rats: ↑ thyroid weight in F1 (both sexes),	Maerkel <i>et al.</i> (2007)
↑ TSH and ↑ T3 in female F1	
Adult oophorectomized rats (12 weeks):	Schmutzler <i>et al.</i> (2007b)
↑ TSH, ↓ T4, ↑ T3,	
↓ Dio 1 activity (kidney),	
ME activity unchanged (liver, kidney)	
Additional organ toxicities and general toxicity	
Developmental study, rats: no effect on body weight in adult F1	Durrer <i>et al.</i> (2005, 2007) and Maerkel <i>et al.</i> (2007)
Developmental study in rats: ↓ thymus weight in adult female F1	Schlumpf <i>et al.</i> (2004b)
Adult oophorectomized rats (3 months): ↑ bone density	Seidlova-Wuttke <i>et al.</i> (2006)
↑ VTG induction and ER α gene expression in liver of male medaka (7 days)	Inui <i>et al.</i> (2003)

Table 2 Continued

Endpoint	References
OMC	
Oestrogen activity: In vitro	
No binding to hER α	Morohoshi et al. (2005)
↑ Transactivation: hER α	Gomez et al. (2005) and Schreurs et al. (2002)
No transactivation: hER α , hER β , rtER α	Kunz et al. (2006), Kunz & Fent (2006b), Morohoshi et al. (2005) and Schreurs et al. (2002)
Antagonism of hER α transactivation	Kunz & Fent (2006b) and Morohoshi et al. (2005)
↑ MCF-7 cell proliferation	Schlumpf et al. (2001b)
Oestrogen activity: Acute in vivo models	
Uterotrophic effect in immature rats	Schlumpf et al. (2001b)
Uterus of adult oophorectomized rats: ↑weight, ↑ ER β and C3 expression	Klammer et al. (2005)
↑VTG induction in male medaka	Inui et al. (2003)
VTG induction in juvenile fathead minnows: no effect	Kunz et al. (2006)
Androgen activity: In vitro	
Antagonism > agonism of hAR transactivation	Kunz & Fent (2006b)
No effect on hAR transactivation	Ma et al. (2003) and Schreurs et al. (2005)
Progesterone activity: In vitro	
Antagonism of hPR transactivation	Schreurs et al. (2005)
Reproductive organs: Developmental toxicity	
No effect on puberty in rats,	Axelstad et al. (2011)
↓ P-testosterone in F1 on PND 16,	
↓prostate weight PND 16 in adult F1,	
altered prostate histology PND 16 in adult F1,	
↓testis weight PND 16, unchanged in adult F1,	
↓ epididymal sperm count adult F1, no effect on uterus or ovary weight	
Reproductive organs: Long-term exposure of adult animals	
Adult oophorectomized rats: uterine weight unchanged or slightly ↑,	Seidlova-Wuttke et al. (2006)
↑ thickness of uterus epithelium, endometrium and myometrium, and of vagina epithelium,	
PR and IGF-1 expression in uterus and vagina	

Table 2 Continued

Endpoint	References
OMC	
Thyroid axis	
In vitro:	Schmutzler et al. (2007b)
↓ iodide uptake,	
↑ TR transactivation (high concentrations)	
Adult oophorectomized rats (5 days):	Klammer et al. (2007)
↓ TSH, ↓ T4, ↓ T3,	
↑ TSH receptor protein, TRH expression in hypothalamus unchanged,	
↓Dio 1 activity (liver)	
Adult oophorectomized rats (3 month):	Schmutzler et al. (2004, 2007b)
↓ T4; T3 and TSH unchanged,	
↓ Dio1 activity (liver, kidney),	
↑ ME activity (kidney, T3 target)	
Developmental study, rats:	Axelstad et al. (2011)
thyroid weight: ↑ in F1 PND 16, unchanged in adult F1,	
T4: ↓ in male F1 PND 16 and in dams, unchanged in female F1 PND 16 and in adult F1.	
Central nervous system: Developmental toxicity	
Developmental study, rats (gavage):	Axelstad et al. (2011)
↓ motor activity in adult female F1,	
↑ spatial learning in adult male F1	
Additional organ toxicities and general toxicity	
Developmental 2 generations study:	Schneider et al. (2005)
↓ body weight gain and ↓ adult body weight in F1 male and female rats (high dose),	
↑ liver weight in female F1	
Developmental study, rats (gavage):	Axelstad et al. (2011)
↓ birth weight and body weight gain, body weight of adult F1: males ↓, females normalized	
Adult oophorectomized rats (5 days):	Klammer et al. (2005)
↓ serum cholesterol, ↓LDL,	
↓ triglycerides and ↓ IGF-1 expression in liver (highest dose)	
OD-PABA	
Oestrogen activity: In vitro	
↑ transactivation: hER α	Gomez et al. (2005) and Schreurs et al. (2002)
↑ transactivation: hER α > hER β ,	Schreurs et al. (2005)
No agonistic action: hER α , rtER α , hER β	Kunz et al. (2006), Kunz & Fent (2006b), Morohoshi et al. (2005) and Schreurs et al. (2002)

Table 2 *Continued*

Endpoint	References
OD-PABA	
Antagonism of hER α transactivation	Kunz & Fent (2006b) and Morohoshi <i>et al.</i> (2005)
No antagonism: hER α or hER β	Schreurs <i>et al.</i> (2002)
↑ MCF-7 cell proliferation	Schlumpf <i>et al.</i> (2001b)
Oestrogen activity: Acute in vivo models	
No uterotrophic effect in immature rats	Schlumpf <i>et al.</i> (2001b)
Androgen activity: In vitro	
Antagonism on hAR transactivation	Kunz & Fent (2006b)
No antagonism on hAR transactivation	Ma <i>et al.</i> (2003)
No agonistic action on hAR transactivation	Kunz & Fent (2006b), Ma <i>et al.</i> (2003) and Schreurs <i>et al.</i> (2005)
Progesterone activity: In vitro	
No agonism or antagonism on hPR transactivation	Schreurs <i>et al.</i> (2005)
PABA	
Oestrogen activity: In vitro	
No binding to hER α	Morohoshi <i>et al.</i> (2005)
Antagonism of hER α transactivation	Kunz & Fent (2006b)
No antagonistic action on hER α	Morohoshi <i>et al.</i> (2005)
No agonistic actions on hER α , rER α	Kunz <i>et al.</i> (2006), Kunz & Fent (2006b) and Morohoshi <i>et al.</i> (2005)
Oestrogen activity: Acute in vivo models	
No data	
Androgen activity: In vitro	
No agonistic or antagonistic activity on hAR, yeast cells	Kunz & Fent (2006b)

oestrogenic in acute in vivo models (Schlumpf *et al.*, 2001b; Schreurs *et al.*, 2002).

Binding affinity to oestrogen receptor α (ER α) and to oestrogen receptor β (ER β) has also been examined (Mueller *et al.*, 2003; Schlumpf *et al.*, 2004a; Morohoshi *et al.*, 2005). The studies differed with respect to ER α and ER β binding preference of individual compounds, but indicate an interaction at the level of ERs. This was also demonstrated by the fact that the proliferative effect of 4-MBC on MCF-7 cells was abolished by the selective ER antagonist ICI182780 (Schlumpf *et al.*, 2001b). Additional effects on oestrogen synthesis, degradation, protein binding, receptor synthesis, etc. cannot be excluded, but have not been investigated as yet.

Interestingly, Kunz and Fent detected antagonistic activity of almost all tested UV-filters in yeast expressing human ER α (hER α): BP-3, 3-BC, 4-MBC, OMC, HMS, OD-PABA and PABA (Kunz & Fent, 2006b). Antioestrogenic activity of 4-MBC, OD-PABA and PABA was supported in a couple of other studies (Mueller *et al.*, 2003;

Morohoshi *et al.*, 2005). In contrast, BP-3, found to be the most antioestrogenic UV-filter in the Kunz and Fent study (Kunz & Fent, 2006b) had in another study only weak binding affinity to hER α , compared with 17 β -estradiol (Morohoshi *et al.*, 2005). Moreover, Schreurs and colleagues also investigated antagonistic oestrogenic activity of BP-3, 3-BC, 4-MBC, HMS, OMC and OD-PABA, but in contrast, did not report any effects of the tested compounds (Schreurs *et al.*, 2002).

These conflicting results regarding antioestrogenic activity are compatible with data from oestrogen agonist studies indicating that the agonistic activity of many UV-filters is of the partial agonist type (Schlumpf *et al.*, 2001b, 2004a). In addition, those studies differ in their type of assay, being also different in their capability to discriminate between agonistic and antagonistic effects.

The oestrogenic activity of BP-3, 3-BC, 4-MBC and OMC was confirmed by acute in vivo tests using increased uterine weight in immature rats (Schlumpf *et al.*, 2001b, 2004a; Tinwell *et al.*, 2002) or oophorectomized rats (Klammer *et al.*, 2005) as endpoint. Furthermore, elevated vitellogenin in fish, a phenotypic endpoint for the oestrogenic action, has been observed in a number of ecotoxicological studies of 3-BC, 4-MBC and OMC (Holbech *et al.*, 2002; Inui *et al.*, 2003; Kunz *et al.*, 2006). However, increased uterine weight following BP-3 exposure of immature rats conflicted with unchanged uterine weight in an adult oophorectomized rat model, indicating possible higher sensitivity of immature rats to BP-3 (Schlumpf *et al.*, 2001b; Schlecht *et al.*, 2004). In vitro oestrogenic activity of HMS and OD-PABA could not be confirmed in vivo as reported (Schlumpf *et al.*, 2001b; Schreurs *et al.*, 2002).

Influence on androgen activity

BP-3, 3-BC, 4-MBC, HMS, OMC and OD-PABA exhibited antiandrogenic activity in vitro, even though data on individual compounds were conflicting (Ma *et al.*, 2003; Schreurs *et al.*, 2005; Kunz & Fent, 2006b). In contrast to other UV-filters, which were mainly androgen antagonists, HMS exhibited both full agonistic and antagonistic androgen activity in vitro by producing full dose-response curve binding to the human androgen receptor (hAR) and inhibiting the action of dihydrotestosterone (DHT) (Kunz & Fent, 2006b). In addition to in vitro antiandrogenic activity, OMC caused a decrease in serum-testosterone among immature offspring in a developmental study in rats (Axelstad *et al.*, 2011). For the remaining compounds in Table 1, antiandrogenic activity of UV-filters has not yet been investigated in vivo.

Influence on progesterone activity

3-BC, 4-MBC, BP-3, OMC and HMS were all tested using a progesterone receptor (PR) CALUX bioassay and all

found to exhibit antagonistic action on the PR in U2-OS cells (Schreurs *et al.*, 2005). The action of those UV-filters could be reversed by the PR agonist ORG2058, indicating a PR mediated action. OD-PABA was also tested, but did not exhibit progesterone activity in vitro (Schreurs *et al.*, 2005).

OMC and 4-MBC were examined for progesterone effect in vivo (Seidlova-Wuttke *et al.*, 2006; Axelstad *et al.*, 2011). Only exposure to OMC in vivo confirmed progesterone activity observed in vitro, resulting in a decrease in the plasma-progesterone concentration in a developmental study in rats (Axelstad *et al.*, 2011) and in altered transcription of PR in uterus and vagina among oophorectomized Sprague–Dawley rats orally exposed for 3 months (Seidlova-Wuttke *et al.*, 2006). In vitro progesterone activity of 4-MBC was not confirmed in the later study performed on oophorectomized rats (Seidlova-Wuttke *et al.*, 2006). This finding does not rule out the possibility of 4-MBC's interference with progesterone signalling in vivo among immature rats, if their sensitivity is higher compared with oophorectomized rats, just as it was the case for oestrogenic activity of BP-3 (Schlumpf *et al.*, 2001b; Schlecht *et al.*, 2004).

Effects on reproductive organs and development

Studies on reproductive and developmental toxicity have been published for only three of the endocrine active UV-filters, namely 4-MBC, 3-BC and OMC. Delay of male puberty and reduced prostate weight were the most sensitive variables for reproductive toxicity following exposure to 3-BC and 4-MBC in extended one generation developmental studies, where Long Evans rats were orally exposed for 10 weeks before mating, during pregnancy and lactation and then their offspring continued oral exposure until adulthood (Schlumpf *et al.*, 2004b; Durrer *et al.*, 2007). Those effects were seen at a dose of 0.24 mg/kg bw/day for 3-BC and of 7 mg/kg bw/day for 4-MBC. In contrast, the reproductive toxicity two generation study with OMC reported only delayed male and female puberty at the highest dose of 1000 mg/kg bw/day, which was not attributed to the compound, but rather to a natural variation within the 'historical control range', in spite of the statistically significant difference compared with control animals in the study (Schneider *et al.*, 2005). Axelstad *et al.* (2011) did not find any effect of OMC on the time of puberty in a one generation developmental study either.

Several other adverse effects on the reproductive system were observed in extended one generation developmental studies after exposure to BP-3, 3-BC, 4-MBC and OMC comprising alteration in weight and histology of reproductive organs in both sexes (Schlumpf *et al.*, 2004b; Durrer *et al.*, 2007; Hofkamp *et al.*, 2008; Axelstad *et al.*, 2011). Developmental studies with 3-BC and 4-MBC (Schlumpf

et al., 2004b; Durrer *et al.*, 2005, 2007) and acute and long-term BP-3 and OMC studies in adult oophorectomized rats (Klammer *et al.*, 2005; Schlecht *et al.*, 2004; Seidlova-Wuttke *et al.*, 2006) found alterations in proteins and gene expression of ER, AR, PR, insulin-like growth factor-I (IGF-1), complement protein 3 (C3), nuclear receptor co-repressor (N-Cor), steroid receptor coactivator-1 (SRC-1) in uterus and prostate. Those findings indicate possible mechanisms of action behind the reproductive toxicity. Alterations in oestrogen target gene expression following peri- and postnatal exposure to 3-BC and 4-MBC also occurred in brain regions important for rats' sexual behaviour (Ventromedial Hypothalamic nucleus (VMH) and Medial Preoptic area (MPO)) (Maerkel *et al.*, 2005, 2007; Faass *et al.*, 2009). This was supported by observed changes in female sexual behaviour, such as reduction in proceptive behaviour, altered attractive behaviour resulting in a decreased number of mounts, impaired receptive behaviour and episodes of rejection following exposure to 3-BC and 4-MBC in an extended one generation developmental study (Faass *et al.*, 2009).

In addition, a 90-day BP-3 study in adult mice (French, 1992) and a 3-BC extended one generation developmental study in rats (Faass *et al.*, 2009) resulted in changes in the oestrous cycle.

Fertility in males was affected in a 90-day study with BP-3, where sperm density decreased in a dose-related manner following dermal exposure in mice and at the highest dose following oral exposure in mice and rats (French, 1992). In addition, at the same dose level an increased number of abnormal spermatozoa was observed in mice. Perinatal and early postnatal exposure to OMC in rats also resulted in decreased sperm count (Schneider *et al.*, 2005; Axelstad *et al.*, 2011).

Reduction of litter size and survival rate in offspring were seen after exposure of dams during pregnancy to higher doses of 3-BC (above 2.4 mg/kg bw/day) and 4-MBC (above 24 mg/kg bw/day) (Schlumpf *et al.*, 2001a, 2004b). The mechanisms behind this perinatal toxicity have not been clarified, but involvement of the immune system and metabolism of the compounds are suspected because the same doses of 4-MBC caused decrease in thymus weight of offspring and increase in weight of thyroid in dams (Schlumpf *et al.*, 2004b).

Effects on hypothalamic–pituitary–thyroid axis

A wide range of in vitro and in vivo studies support that 4-MBC, BP-3 and OMC may interfere with the hypothalamic–pituitary–thyroid axis (HPT).

An in vitro and a 5-day in vivo study with BP-3 have shown that this compound interacts with thyroid function by an agonistic effect on the thyroid receptor (TR) in HepG2 cells (Schmutzler *et al.*, 2007b) and by decreasing

the expression of the *ERR1* gene in the thyroid gland (Schlecht *et al.*, 2004). Whether those findings result in any adverse effect on the thyroid axis have not been examined and further investigations on BP-3 in long-term studies seem indicated.

An adverse effect on the thyroid axis indicated by alterations in the concentrations of thyroid hormones following exposure to 4-MBC and OMC was found in 90 days toxicological studies (Schmutzler *et al.*, 2004, 2007b). An extended one generation developmental study in rats confirmed alterations in thyroid-stimulating hormone (TSH) and total triiodothyronine (T3) following 4-MBC exposure, supplemented with increased thyroid weight in offspring (Maerkel *et al.*, 2007). A developmental study of OMC resulted in decreased total thyroxine (T4) in dams and in juvenile male offspring and in increased weight of thyroid gland in juvenile rats of both sexes (Axelstad *et al.*, 2011). A sex difference was noted: female offspring were less sensitive to OMC exposure, resulting in unaltered T4 (Axelstad *et al.*, 2011).

The mechanisms behind the adverse effects observed in the thyroid axis following *in vivo* exposure to 4-MBC and OMC could be partially explained by a decrease in Dio 1 activity, an enzyme promoting both activation and inactivation of thyroid hormones and by decreased iodide uptake in FRTL-5 cells (Klammer *et al.*, 2007; Schmutzler *et al.*, 2007b). In addition, OMC exhibited agonistic action on TR in the HepG2 cell line (Schmutzler *et al.*, 2007b). In contrast, thyroid peroxidase (TPO) activity was not affected neither following OMC exposure nor 4-MBC exposure (Schmutzler *et al.*, 2004; Klammer *et al.*, 2007).

The UV-filter benzophenone-2 (BP-2) is not allowed to be used in cosmetics in EU (European Commission), but is still used in USA to protect cosmetic products against UV-rays (Food and Drug Administration [FDA], 2012). This is of concern as it has been shown to disturb thyroid function in an acute toxicity study on oophorectomized Sprague Dawley rats, altering TSH, T4 and Dio 1 activity and to decrease human receptor TPO (hrTPO) activity *in vitro* (Schmutzler *et al.*, 2007a).

Several UV-filters, including 3-BC, HMS, OD-PABA or PABA seem not to have been examined for their possible effects on the thyroid axis.

General toxicity

In mammalian long-term exposure models, general toxicity evaluated by alterations of food consumption and in body and liver weights was found in the higher dose range after exposure to BP-3 (French, 1992), 3-BC (>2.4 mg/kg bw/day) (Schlumpf *et al.*, 2004b) and OMC (>500mg/kg bw/day) (Schneider *et al.*, 2005; Axelstad *et al.*, 2011). Liver and kidney weights were affected after both dermal and oral exposure to BP-3 (French, 1992).

In contrast, histological alterations in liver and kidney were observed only after oral exposure to BP-3 in the latter study. 4-MBC reduced body weight only transiently in postnatal 4-MBC exposed F1 offspring. Body weights were again normal at puberty and in adulthood also in the higher dose groups, and no signs of general toxicity were observed in the parent animals (F0) (Durrer *et al.*, 2007; Maerkel *et al.*, 2007). Effects on reproductive organs and development were also present at doses devoid of general toxicity. Indications of general toxicity were further observed in experiments on fish exposed to 3-BC and 4-MBC, where body weight decreased in a dose-dependent manner (Kunz *et al.*, 2006). A few deaths were observed by Schneider and colleagues in a two generation study in rats, but were not considered to be related to OMC exposure (Schneider *et al.*, 2005).

Human exposure to sunscreens

Table 3 summarizes prevailing data on human exposure to chemical UV-filters used in cosmetics. Experimental studies showed that BP-3, 4-MBC and OMC rapidly permeated intact skin (Gustavsson *et al.*, 2002; Janjua *et al.*, 2004, 2008; Gonzalez *et al.*, 2006) and could be detected in plasma after 1–2 h following application (Fig. 2) (Janjua *et al.*, 2008). Interestingly, the concentrations of these compounds in the same experimental study in male urine and plasma were higher than in female samples (Janjua *et al.*, 2004), indicating a gender difference in the metabolism, distribution and possibly also in the accumulation of UV-filters in adipose tissue.

Furthermore, Table 3 shows a substantial exposure of the general population to UV-filters. BP-3, the most common UV-filter in the USA, was found in more than 96% of 2517 urine samples collected throughout 1-year (2003–2004) from the general US population in the NHANES study (Calafat *et al.*, 2008). BP-3 was also detected in all urine samples collected from 129 Danish children and adolescents in the month of November, even though days are short and sun protection is not needed at that time of year (H. Frederiksen, O. Nielsen, L. Aksglaede, K. Sorensen, T. H. Lassen, N. E. Skakkebaek, K. Main, A. Juul & A. Andersson, unpublished data, 2012).

Breastfed babies are exposed to UV-filters through breast milk (Schlumpf *et al.*, 2010). One or more UV-filters were present in 85% of Swiss human milk samples (Schlumpf *et al.*, 2010). Bisphenol A with a similar chemical structure to BP-3, were shown to pass the blood-placenta barrier (Schonfelder *et al.*, 2002; Lee *et al.*, 2008). Thus in theory, chemicals like BP-3 may also pass the blood-placenta barrier. Studies investigating amniotic fluid are required to investigate whether perinatal exposure to UV-filters, which are found in the urine of pregnant women occurs (Wolff

Table 3 Human exposure to UV-filters

Study design	Number of test subjects	Sample type	UV-filter	Reported use(%)	Positive ^d samples(%)	Concentrations	Ref.
Observational studies							
USA: NHANES study 2003–2004 based on US general population > 6 years of age	2517 people	Urine	BP-3	–	96.8	22.9 (18.1–28.9) µg/L ^c	Calafat et al. (2008)
USA: The Children's Environmental Health study 1998–2002: Multiethnic prospective cohort of pregnant women-infant pairs in New York during 3. trimester	404 pregnant women	Urine	BP-3	–	97.8	7.5 µg/L ^a	Wolff et al. (2008)
France: Eden Mother-Child cohort recruited before gestational week 28 in 2003–2006	191 pregnant women	Urine	BP-3	–	80.5	1.3 µg/L ^a	Philippat et al. (2012)
Switzerland: Cohorts 2004–2006 Mothers who gave birth to a single child at the University Women's Hospital Basel	54 women	Human breast milk	BP-3 4-MBC OMC HMS OCT OD-PABA 3-BC	13.21 26.42 66.04 15.09 43.40 1.89 0	12.96 20.37 77.78 5.56 66.67 1.85 0	52.23 ± 50.69 ng/g lipid ^a 22.12 ± 12.80 ng/g lipid ^a 27.50 ± 22.15 ng/g lipid ^a 29.37 ± 27.64 ng/g lipid ^a 30.18 ± 24.51 ng/g lipid ^a 49.00 ng/g lipid ^c	Schlumpf et al. (2010)
Experimental studies							
Denmark: Single blinded experimental study 1 week with UV-filters free lotion + 1 week with Daily whole-body application of sunscreen 2 mg/cm ² with BP-3, 4-MBC and OMC Concentration 10% of each	32: 15 males + 17 postmenopausal females	Plasma	BP-3 4-MBC OMC BP-3 4-MBC OMC	100	100	Female (ng/ml) 200 ^b 20 ^b 10 ^b 60 ^b 5 ^b 5 ^b Male (ng/ml) 300 ^b 20 ^b 20 ^b 140 ^b 7 ^b 8 ^b	Janjua et al. (2004)
Sweden: Experimental study 1x whole-body application of sunscreen with BP-3 2mg/cm ² Concentration 4%	11: 7 males + 4 females	Plasma	BP-3 4-MBC OMC	100	100	187 ^b 16 ^b 7 ^b	Janjua et al. (2008)
Sweden: Experimental study: hole-body application of 2 mg/cm ² sunscreen SPF14 with 4% BP-3 twice a day for 5 days One half of participants was daily irradiated with UVA+UVB according to Fitzpatrick skin type	25: 16 women and 9 men	Urine	BP-3 4-MBC OMC	100	100	44 ^b 4 ^b 6 ^b 9.8 mg = 0.5% of the applied amount	Gustavsson et al. (2002)
		Urine:	BP-3	100	100	3.7% of applied amount	Gonzalez et al. (2006)

^aMedian concentration.^bMaximum median concentration.^cGeometric mean concentration.^dPercentage of samples with content of UV-filters above analytical detection limit.




Figure 2 Plasma concentration of UV-filters BP-3, 4-MBC and OMC among males after one dermal application daily for four days (Janjua *et al.*, 2008).

et al., 2008). High concentrations of BP-3 in mothers' urine were associated with decreased birth weight in girls and increased birth weight and head circumference in boys (Wolff *et al.*, 2008; Philippat *et al.*, 2012). Human studies with genital malformations in infants as an endocrine specific endpoint are required to clarify a possible endocrine disrupting effect of UV-filters on the human foetus. However, developmental animal studies where rats were exposed to UV-filters, did not report any reduction of the anogenital distance (ADG) or increased rate of genital malformations.

In spite of the wide human exposure to UV-filters only few studies have examined the effects of UV-filters on humans (Janjua *et al.*, 2004, 2007; Jannesson *et al.*, 2004; Wolff *et al.*, 2008; Philippat *et al.*, 2012). A double blinded clinical trial measuring the gingival index, that relate to the severity and location of periodontal disease, showed that dentifrice containing BP-3 reduced periodontal disease by 25% (Jannesson *et al.*, 2004), which supported the *in vitro* data suggesting BP-3 to be an inhibitor of PG synthesis (Jannesson *et al.*, 2004; Kristensen *et al.*, 2011).

Janjua and colleagues examined the effects on reproductive (Janjua *et al.*, 2004) and thyroid (Janjua *et al.*, 2007) hormones following dermal application of a mixture of BP-3, 4-MBC and OMC. A significant increase in inhibin B and a decrease in free triiodothyronine (FT3), free thyroxine (FT4), T3, T4, thyroxine-binding globulin (TBG) and testosterone were seen, but were not considered to be related to the application of a sunscreen mixture, but rather to biological variation (Janjua *et al.*, 2007). However, the duration of those studies was too short to be conclusive.

Discussion

As summarized in this review, a large number of *in vivo* animal studies and *in vitro* studies have shown that there are numerous potential adverse effects of UV-filters pres-

ent in sunscreens and cosmetics. The effects include developmental and reproductive effects, apparently caused by endocrine disrupting actions of these chemicals. Other studies could not find such adverse effects. However, because of the wide human exposure in combination with the clear endocrine disruptive effects observed in a large number of well designed studies, the UV-filters BP-3, 4-MBC and OMC can be considered as substances of high concern in relation to human risk.

Importantly, most of the studied adverse effects of UV-filters have been evaluated after oral exposure. However, the primary exposure of humans to UV-filters via cosmetics occurs through dermal application. Therefore, the UV-filters enter the systemic circulation directly without first being metabolized by passage through the liver, thereby leading to a greater risk of the compounds reaching all tissues of the body unaltered, as was observed in rats following dermal exposure to 3-BC (Søeborg *et al.*, 2006). In addition, a three-fold greater oestrogenic effect of 4-MBC in rats was observed after topical application compared with oral exposure indicating higher bioavailability of the compound (Schlumpf *et al.*, 2001b).

Another challenge in studies of sunscreens in cosmetics is that the products often contain several UV-filters in combination. The total effect of these mixtures are poorly examined although a few existing studies have shown that mixtures of chemicals, including UV-filters, might act additively and exhibit toxic activity, even at the no observed adverse effect level (NOAEL) of the individual compounds (Heneweer *et al.*, 2005; Kunz & Fent, 2006a; Kortenkamp *et al.*, 2007).

Humans are not only exposed to UV-filters when the agents are used for sun protection of the skin. Exposure apparently occurs from various sources. Almost all samples in NHANES study (Calafat *et al.*, 2008) and all samples in a Danish children cohort (H. Frederiksen, O. Nielsen, L. Aksglaede, K. Sorensen, T. H. Lassen, N. E. Skakkebaek, K. Main, A. Juul & A. Andersson, unpublished data, 2012), contained BP-3, indicating year round exposure independent of sunscreen use. The presence of UV-filters in milk of Swiss mothers was correlated with use of sunscreens in 55% of the cases; in 60% of the cases, the presence of these compounds in milk was related to the use of other cosmetic products containing UV-filters (Schlumpf *et al.*, 2010). The likely sources may be hair spray, lipsticks, shampoo, make-up, perfumes, skin care products as well as non-cosmetic products, such as carpets, furniture, clothing and washing powder (Schlecht *et al.*, 2004; Morohoshi *et al.*, 2005; Kunz & Fent, 2006b; Schlumpf *et al.*, 2010). Here, the UV-filters are used to protect the products from effects of UV-radiation. Considering these findings, it cannot be ruled out that a considerable part of the total human exposure to UV-filters might occur via products other than sunscreens.

It is of particular concern that human babies are exposed to UV-filters through breast milk (Schlumpf *et al.*, 2010). The highest concentration of 4-MBC found in human milk was 48.37 ng/g lipid (Schlumpf *et al.*, 2010), which was only 4.3 times lower than the concentration of 4-MBC in rat milk (208.6 ng/g lipid) (Schlumpf *et al.*, 2008) following oral exposure to 4-MBC at the lowest observed adverse effect level (LOAEL) (7 mg/kg/day), with delay of male puberty and prostate weight as endpoints (Durrer *et al.*, 2007).

In conclusion, it is of concern that (1) a large number of in vitro and in vivo animal studies have shown endocrine disrupting effects of UV-filters present in sunscreens, although other studies failed to find such effects and (2) application of cosmetics with UV-filters to the skin can result in absorption of UV-filters into the human systemic circulation and subsequently might result in exposure of all tissues in the body. Considering these facts together with the wide and increasing use of sunscreens and the increasing incidence of malignant melanoma, for which UV-filters are assumed to protect, it seems pertinent to investigate whether sunscreen use in humans on balance is beneficial for human health.

Acknowledgement

The authors would like to acknowledge ©The Nielsen Company for providing data on annual sunscreen consumption in Denmark 1999–2010. Sreeram V. Ramagopalan is highly acknowledged for providing raw data used in his correspondence 'The questionable effectiveness of sunscreen', A.E. Handel and S.V. Ramagopalan, *The Lancet* (Handel & Ramagopalan, 2010). Furthermore, we would like to thank Claus Jørgensen, Senior Environmental and Transport adviser, Danish consumer council for providing international contacts, useful in our sunscreen use research. Kirsten and Freddy Johansen's Foundation is thanked for their financial support.

References

- Australian Institute of Health and Welfare (AIHW). Age standardise Malignant Melanoma Incidence in Australia. Available at: <http://www.aihw.gov.au/acim-books/> (Last accessed 3 April 2012).
- Axelstad M, Boberg J, Hougaard KS, Christiansen S, Jacobsen PR, Mandrup KR, Nellemann C, Lund SP & Hass U. (2011) Effects of pre- and postnatal exposure to the UV-filter octyl methoxycinnamate (OMC) on the reproductive, auditory and neurological development of rat offspring. *Toxicol Appl Pharmacol* 250, 278–290.
- Calafat AM, Wong LY, Ye X, Reidy JA & Needham LL. (2008) Concentrations of the sunscreen agent benzophenone-3 in residents of the United States: National Health and Nutrition Examination Survey 2003–2004. *Environ Health Perspect* 116, 893–897.
- Cancer Research UK. Age standardise Malignant Melanoma incidence in UK. Available at: <http://info.cancerresearchuk.org/cancerstats/types/skin/incidence/#Trends> (Last accessed 3 April 2012).
- Deep S. (2010) Sunscreen: A History. *New York Times*. Available at: http://www.nytimes.com/2010/06/24/fashion/24skinside.html?_r=1 (Last accessed 3 April 2012).
- Dupuy A, Dunant A & Grob JJ. (2005) Randomized controlled trial testing the impact of high-protection sunscreens on sun-exposure behavior. *Arch Dermatol* 141, 950–956.
- Durrer S, Maerkel K, Schlumpf M & Lichtensteiger W. (2005) Estrogen target gene regulation and coactivator expression in rat uterus after developmental exposure to the ultraviolet filter 4-methylbenzylidene camphor. *Endocrinology* 146, 2130–2139.
- Durrer S, Ehnes C, Fuetsch M, Maerkel K, Schlumpf M & Lichtensteiger W. (2007) Estrogen sensitivity of target genes and expression of nuclear receptor co-regulators in rat prostate after pre- and postnatal exposure to the ultraviolet filter 4-methylbenzylidene camphor. *Environ Health Perspect* 115(Suppl 1), 42–50.
- European Commission. Regulation (EC) No 1223/2009 of the European parliament and of the council of 30 November 2009 on cosmetic products. Annex VI. List of UV-filters allowed in cosmetic products Available at: http://ec.europa.eu/consumers/cosmetics/cosing/index.cfm?fuseaction=search.results&annex_v2=VI&search (Last accessed 3 April 2012).
- European Commission. Definition of endocrine disruptors. Available at: http://ec.europa.eu/research/endocrine/background_disruption_en.html (Last accessed 3 April 2012).
- Faass O, Schlumpf M, Reolon S, Henseler M, Maerkel K, Durrer S & Lichtensteiger W. (2009) Female sexual behavior, estrous cycle and gene expression in sexually dimorphic brain regions after pre- and postnatal exposure to endocrine active UV-filters. *Neurotoxicology* 30, 249–260.
- Food and Drug Administration (FDA). (2012) OCT Ingredient List (By monograph category). www.fda.gov
- French JE. (1992) NTP technical report on the toxicity studies of 2-Hydroxy-4-methoxybenzophenone (CAS No. 131-57-7) Administered Topically and in Dosed Feed to F344/N Rats and B6C3F1 Mice. *Toxic Rep Ser* 21, 1-E14.
- Gandini S, Sera F, Cattaruzza MS, Pasquini P, Picconi O, Boyle P & Melchi CF. (2005) Meta-analysis of risk factors for cutaneous melanoma: II. Sun exposure. *Eur J Cancer* 41, 45–60.
- Gomez E, Pillon A, Fenet H, Rosain D, Duchesne MJ, Nicolas JC, Balaguer P & Casellas C. (2005) Estrogenic activity of cosmetic components in reporter cell lines: Parabens, UV screens, and musks. *J Toxicol Environ Health A* 68, 239–251.
- Gonzalez H, Farbrot A, Larko O & Wennberg AM. (2006) Percutaneous absorption of the sunscreen benzophenone-3 after repeated whole-body applications, with and without ultraviolet irradiation. *Br J Dermatol* 154, 337–340.
- Green A, Williams G, Neale R, Hart V, Leslie D, Parsons P, Marks GC, Gaffney P, Battistutta D, Frost C, Lang C & Russell A. (1999) Daily sunscreen application and betacarotene supplementation in prevention of basal-cell and squamous-cell carcinomas of the skin: A randomised controlled trial. *Lancet* 354, 723–729.
- Green A, Williams G, Logan V & Strutton G (2011) Reduced melanoma after regular sunscreen use: Randomized trial follow-up. *J Clin Oncol* 29, 257–263.
- Gustavsson GH, Farbrot A & Larko O. (2002) Percutaneous absorption of benzophenone-3, a common component of topical sunscreens. *Clin Exp Dermatol* 27, 691–694.
- Handel AE & Ramagopalan SV (2010) The questionable effectiveness of sunscreen. *Lancet* 376, 161–162.

- Heneweer M, Muusse M, Van den BM & Sanderson JT. (2005) Additive estrogenic effects of mixtures of frequently used UV-filters on pS2-gene transcription in MCF-7 cells. *Toxicol Appl Pharmacol* 208, 170–177.
- Hofkamp L, Bradley S, Tresguerres J, Lichtensteiger W, Schlumpf M & Timms B. (2008) Region-specific growth effects in the developing rat prostate following fetal exposure to estrogenic ultraviolet filters. *Environ Health Perspect* 116, 867–872.
- Holbech H, Norum U, Korsgaard B & Poul B. (2002) The chemical UV-filter 3-benzylidene camphor causes an oestrogenic effect in an in vivo fish assay. *Pharmacol Toxicol* 91, 204–208.
- IARC. CI5plus Database. Age Standardised Incidence Rate in DK. Available at: <http://ci5.iarc.fr/CI5plus/ci5plus.htm> (Last accessed 3 April 2012).
- Inui M, Adachi T, Takenaka S et al. (2003) Effect of UV screens and preservatives on vitellogenin and choriogenin production in male medaka (*Oryzias latipes*). *Toxicology* 194, 43–50.
- Janjua NR, Mogensen B, Andersson AM, Petersen JH, Henriksen M, Skakkebaek NE & Wulf HC. (2004) Systemic absorption of the sunscreens benzophenone-3, octyl-methoxycinnamate, and 3-(4-methyl-benzylidene) camphor after whole-body topical application and reproductive hormone levels in humans. *J Invest Dermatol* 123, 57–61.
- Janjua NR, Kongshoj B, Petersen JH & Wulf HC. (2007) Sunscreens and thyroid function in humans after short-term whole-body topical application: A single-blinded study. *Br J Dermatol* 156, 1080–1082.
- Janjua NR, Kongshoj B, Andersson AM & Wulf HC. (2008) Sunscreens in human plasma and urine after repeated whole-body topical application. *J Eur Acad Dermatol Venereol* 22, 456–461.
- Jannesson L, Birkhed D, Scherl D, Gaffar A & Renvert S. (2004) Effect of oxybenzone on PGE2-production in vitro and on plaque and gingivitis in vivo. *J Clin Periodontol* 31, 91–94.
- Klammer H, Schlecht C, Wuttke W & Jarry H. (2005) Multi-organic risk assessment of estrogenic properties of octyl-methoxycinnamate in vivo: A 5-day sub-acute pharmacodynamic study with ovariectomized rats. *Toxicology* 215, 90–96.
- Klammer H, Schlecht C, Wuttke W, Schmutzler C, Gotthardt I, Kohrle J & Jarry H. (2007) Effects of a 5-day treatment with the UV-filter octyl-methoxycinnamate (OMC) on the function of the hypothalamo-pituitary-thyroid function in rats. *Toxicology* 238, 192–199.
- Kortenkamp A, Faust M, Scholze M & Backhaus T. (2007) Low-level exposure to multiple chemicals: Reason for human health concerns? *Environ Health Perspect* 115(Suppl 1), 106–114.
- Kristensen DM, Skalkam ML, Audouze K et al. (2011) Many putative endocrine disruptors inhibit prostaglandin synthesis. *Environ Health Perspect* 119, 534–541.
- Kunz PY & Fent K. (2006a) Estrogenic activity of UV-filter mixtures. *Toxicol Appl Pharmacol* 217, 86–99.
- Kunz PY & Fent K. (2006b) Multiple hormonal activities of UV-filters and comparison of in vivo and in vitro estrogenic activity of ethyl-4-aminobenzoate in fish. *Aquat Toxicol* 79, 305–324.
- Kunz PY, Galicia HF & Fent K. (2006) Comparison of in vitro and in vivo estrogenic activity of UV-filters in fish. *Toxicol Sci* 90, 349–361.
- Lee YJ, Ryu HY, Kim HK et al. (2008) Maternal and fetal exposure to bisphenol A in Korea. *Reprod Toxicol* 25, 413–419.
- Ma R, Cotton B, Lichtensteiger W & Schlumpf M. (2003) UV-filters with antagonistic action at androgen receptors in the MDA-kb2 cell transcriptional-activation assay. *Toxicol Sci* 74, 43–50.
- Maerkel K, Lichtensteiger W, Durrer S, Conscience M & Schlumpf M. (2005) Sex- and region-specific alterations of progesterone receptor mRNA levels and estrogen sensitivity in rat brain following developmental exposure to the estrogenic UV-filter 4-methylbenzylidene camphor. *Environ Toxicol Pharmacol* 19, 761–765.
- Maerkel K, Durrer S, Henseler M, Schlumpf M & Lichtensteiger W. (2007) Sexually dimorphic gene regulation in brain as a target for endocrine disruptors: Developmental exposure of rats to 4-methylbenzylidene camphor. *Toxicol Appl Pharmacol* 218, 152–165.
- Morohoshi K, Yamamoto H, Kamata R, Shiraiishi F, Koda T & Morita M. (2005) Estrogenic activity of 37 components of commercial sunscreen lotions evaluated by in vitro assays. *Toxicol In Vitro* 19, 457–469.
- Mueller SO, Kling M, Arifin FP, Mecky A, Duranti E, Shields-Botella J, Delansorne R, Broschard T & Kramer PJ. (2003) Activation of estrogen receptor alpha and ERbeta by 4-methylbenzylidene-camphor in human and rat cells: Comparison with phyto- and xenoestrogens. *Toxicol Lett* 142, 89–101.
- Philippat C, Mortamais M & Chevrier C et al. (2012) Exposure to phthalates and phenols during pregnancy and offspring size at birth. *Environ Health Perspect* 120, 464–470.
- Rebut D. (1990) The Sunscreen Industry in Europe: Past, present, and future. In: Sunscreens: Development, evaluation and regulatory aspects (ed. D Rebut), pp. 161–171. Marcel Dekker, New York, NY.
- Schlecht C, Klammer H, Jarry H & Wuttke W. (2004) Effects of estradiol, benzophenone-2 and benzophenone-3 on the expression pattern of the estrogen receptors (ER) alpha and beta, the estrogen receptor-related receptor 1 (ERR1) and the aryl hydrocarbon receptor (AhR) in adult ovariectomized rats. *Toxicology* 205, 123–130.
- Schlumpf M, Berger B, Cotton M, Conscience-Egli S, Durrer I, Fleischmann V, Haller K, Maerkel W & Lichtensteiger W. (2001a) Estrogen active UV screens. *SÖFW-J* 127, 10–15.
- Schlumpf M, Cotton B, Conscience M, Haller V, Steinmann B & Lichtensteiger W. (2001b) In vitro and in vivo estrogenicity of UV screens. *Environ Health Perspect* 109, 239–244.
- Schlumpf M, Jarry H, Wuttke W, Ma R & Lichtensteiger W. (2004a) Estrogenic activity and estrogen receptor beta binding of the UV-filter 3-benzylidene camphor. Comparison with 4-methylbenzylidene camphor. *Toxicology* 199, 109–120.
- Schlumpf M, Schmid P, Durrer S et al. (2004b) Endocrine activity and developmental toxicity of cosmetic UV-filters – an update. *Toxicology* 205, 113–122.
- Schlumpf M, Kypke K, Vökt C et al. (2008) Endocrine active UV-filters: Developmental toxicity and exposure through breast milk. *Chimia* 62, 345–351.
- Schlumpf M, Kypke K, Wittassek M, Angerer J, Mascher H, Mascher D, Vökt C, Birchler M & Lichtensteiger W. (2010) Exposure patterns of UV-filters, fragrances, parabens, phthalates, organochlor pesticides, PBDEs, and PCBs in human milk: Correlation of UV-filters with use of cosmetics. *Chemosphere* 81, 1171–1183.
- Schmutzler C, Hamann I, Hofmann PJ et al. (2004) Endocrine active compounds affect thyrotropin and thyroid hormone levels in serum as well as endpoints of thyroid hormone action in liver, heart and kidney. *Toxicology* 205, 95–102.
- Schmutzler C, Bacinski A, Gotthardt I et al. (2007a) The ultraviolet filter benzophenone 2 interferes with the thyroid hormone axis in rats and is a potent in vitro inhibitor of human recombinant thyroid peroxidase. *Endocrinology* 148, 2835–2844.
- Schmutzler C, Gotthardt I, Hofmann PJ et al. (2007b) Endocrine disruptors and the thyroid gland – a combined in vitro and in vivo

- analysis of potential new biomarkers. *Environ Health Perspect* 115(Suppl 1), 77–83.
- Schneider S, Deckardt K, Hellwig J, Kuttler K, Mellert W, Schulte S & van RB. (2005) Octyl methoxycinnamate: Two generation reproduction toxicity in Wistar rats by dietary administration. *Food Chem Toxicol* 43, 1083–1092.
- Schonfelder G, Wittfoht W, Hopp H, Talsness CE, Paul M & Chahoud I. (2002) Parent bisphenol A accumulation in the human maternal–fetal–placental unit. *Environ Health Perspect* 110, A703–A707.
- Schreurs R, Lanser P, Seinen W & van der BB. (2002) Estrogenic activity of UV-filters determined by an in vitro reporter gene assay and an in vivo transgenic Zebrafish assay. *Arch Toxicol* 76, 257–261.
- Schreurs RH, Sonneveld E, Jansen JH, Seinen W & van der BB. (2005) Interaction of polycyclic musks and UV-filters with the estrogen receptor (ER), androgen receptor (AR), and progesterone receptor (PR) in reporter gene bioassays. *Toxicol Sci* 83, 264–272.
- SEER. Age standardise Malignant Melanoma Incidence in USA. Available at: <http://seer.cancer.gov/faststats/selections.php?#Output> (Last accessed 3 April 2012).
- Seidlova-Wuttke D, Jarry H, Christoffel J, Rimoldi G & Wuttke W. (2006) Comparison of effects of estradiol (E2) with those of octyl-methoxycinnamate (OMC) and 4-methylbenzylidene camphor (4MBC) – 2 filters of UV light – on several uterine, vaginal and bone parameters. *Toxicol Appl Pharmacol* 210, 246–254.
- Søeborg T, Ganderup NC, Kristensen JH, Bjerregaard P, Pedersen KL, Bollen P, Hansen SH & Halling-Sorensen B. (2006) Distribution of the UV-filter 3-benzylidene camphor in rat following topical application. *J Chromatogr B Analyt Technol Biomed Life Sci* 834, 117–121.
- Sundhedsstyrelsen. (2012) Age standardise Malignant Melanoma incidence in DK 2001–2010. Cancerregisteret 2010
- Thompson SC, Jolley D & Marks R. (1993) Reduction of solar keratoses by regular sunscreen use. *N Engl J Med* 329, 1147–1151.
- Tinwell H, Lefevre PA, Moffat GJ, Burns A, Odum J, Spurway TD, Orphanides G & Ashby J. (2002) Confirmation of uterotrophic activity of 3-(4-methylbenzylidene)camphor in the immature rat. *Environ Health Perspect* 110, 533–536.
- Wang SQ, Setlow R, Berwick M, Polsky D, Marghoob AA, Kopf AW & Bart RS. (2001) Ultraviolet A and melanoma: A review. *J Am Acad Dermatol* 44, 837–846.
- Wolff MS, Engel SM, Berkowitz GS, Ye X, Silva MJ, Zhu C, Wetmur J & Calafat AM. (2008) Prenatal phenol and phthalate exposures and birth outcomes. *Environ Health Perspect* 116, 1092–1097.

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Figure S1. Purchase of sunscreen products by volume per head over time in the US, UK, AUS and DK.

Table S1. In vitro and in vivo effects of UV-filters in animals (Presence of effect: +; Absence of effect: –; Increased: ↑; Decreased: ↓).

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.